

Dust-shell Universe in the modified gravity scenario

Michael Maziashvili*

*Department of Theoretical Physics, Tbilisi State University,
3 Chavchavadze Ave., Tbilisi 0128, Georgia*

The dynamics of the dust-shell model of universe is exactly solved for the modified Schwarzschild solution. This solution is used to derive the cosmology corresponding to the modified gravity.

PACS numbers: 04.50.+h, 95.30.Sf, 98.80.Jk

I. INTRODUCTION

The CMBR gives a strong observational basis for the assumption of homogeneity and isotropy of our universe. Indeed, the deviation of universe from a homogenous and isotropic space is $\sim 10^{-5}$ at the stage of decoupling. Thus our universe is well approximated by the Friedmann-Lemaître model before this stage. On the other hand the present universe is highly inhomogeneous on small scales. The model consisting of the spherical shells of dust with the common center, where the rest observer is situated, was considered in [1, 2] to study how the small scale inhomogeneities affect the global dynamics when averaged on larger scales. The gravitational field in the region between the shells is described due to Birkhoff's theorem by the standard Schwarzschild geometry and Israel's [3] matching conditions are used to derive the equations of motion of the dust shell. The dynamics of this model is exactly integrable that allows one to make the analytical evaluations. On the other hand in the last two decades Milgrom's modified Newtonian dynamics (MOND) paradigm [4] has gained recognition as a successful scheme for unifying much of extragalactic dynamics phenomenology without invoking dark matter [5]. By presuming the Birkhoff's law the relation between modified gravity and corresponding cosmology was established in [6]. In this framework the viability of MOND cosmology was considered in [7]. In this brief note we consider the dust shell universe under assumption of modified Schwarzschild solution. Throughout this paper we put $G = c = 1$.

Let us begin by considering of a single dust shell under assumption of modified Schwarzschild solution given by

$$ds^2 = f^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) - f dt^2, \quad (1)$$

where $f = 1 - 2g(m/r)$ with some function g differing from the standard Schwarzschild solution $g(x) = x$. Using Israel's matching conditions [3] the equation of motion of a spherical shell of dust takes the form

$$\ddot{R} \left\{ \sqrt{1 + \dot{R}^2} + \sqrt{1 + \dot{R}^2 - 2g\left(\frac{m}{R}\right)} \right\} =$$

$$- g' \left(\frac{m}{R} \right) \frac{m \sqrt{1 + \dot{R}^2}}{R^2}, \quad (2)$$

where the prime stands for the derivative with respect to the whole argument and over-dot denotes derivative with respect to the proper time τ . The relation between t and τ is

$$\frac{dt}{d\tau} = [1 - 2g(m/R)]^{-1} \left[1 - 2g\left(\frac{m}{R}\right) + \left(\frac{dR}{d\tau}\right)^2 \right]. \quad (3)$$

The first integral of eq.(2) is given by

$$(1 + \dot{R}^2)^{1/2} = a + \frac{g(m/R)}{2a}, \quad (4)$$

where a is the constant of integration characterizing the binding energy of the shell. From eq.(4) one sees that in the case of standard Schwarzschild solution expanding shell with positive binding energy corresponding to $a < 1$ falls back after reaching a finite maximal radius. In the case of MOND the gravity at large distances behaves as [7]

$$g\left(\frac{m}{r}\right) \sim \ln\left(\frac{2m}{r^3 H_0^2}\right) + const.,$$

or if accommodated to the cosmological data [7]

$$g\left(\frac{m}{r}\right) \sim (r H_0)^2,$$

where H_0 denotes the present value of the Hubble constant. So, in the case of MOND the dust shell can tunnel through the potential barrier to the regime of unbounded expansion.

Let us now consider the dust shell universe which represents the spherically symmetric shells of dust with a common center where the rest observer is situated. Denoting the mass and radius of the i th shell by m_i and R_i the equation of motion of the i th shell takes the form

$$\ddot{R}_i \left\{ \left[1 + \dot{R}_i^2 - 2g\left(\frac{m_{i-1}}{R_i}\right) \right]^{1/2} + \left[1 + \dot{R}_i^2 - 2g\left(\frac{m_i}{R_i}\right) \right]^{1/2} \right\} =$$

*Electronic address: maziashvili@ictsu.tsu.edu.ge

$$-\frac{m_{i-1}g'(m_{i-1}/R_i)}{R_i^2} \left[1 + \dot{R}_i^2 - 2g\left(\frac{m_i}{R_i}\right) \right]^{1/2} - \frac{m_i g'(m_i/R_i)}{R_i^2} \left[1 + \dot{R}_i^2 - 2g\left(\frac{m_{i-1}}{R_i}\right) \right]^{1/2}. \quad (5)$$

The first integral of this equation can be written as

$$1 + \dot{R}_i^2 = a_i^2 + g\left(\frac{m_{i-1}}{R_i}\right) + g\left(\frac{m_i}{R_i}\right) + \frac{1}{4a_i^2} \left[g\left(\frac{m_i}{R_i}\right) - g\left(\frac{m_{i-1}}{R_i}\right) \right]^2. \quad (6)$$

Recalling that in the standard case the matter density ρ_i is defined as [1, 2]

$$\frac{m_{i-1} + m_i}{2} = \frac{4}{3}\pi\rho_i x_i^3,$$

where x_i denotes the initial radius of the i th shell, we find it natural to define the matter density in the case of modified gravity as follows

$$\frac{1}{2} \left[g\left(\frac{m_{i-1}}{R}\right) + g\left(\frac{m_i}{R}\right) \right] = g\left(\frac{4}{3}\pi\rho_i \frac{x_i^3}{R}\right).$$

If one sets the parameters containing in eq.(6) as follows

$$\rho_i = \rho, \quad a_i^2 - 1 = -kx_i^2$$

the expansion law of the dust shell can be written as

$$\left(\frac{\dot{R}_i}{R_i}\right)^2 = -k\left(\frac{x_i}{R_i}\right)^2 + \frac{2}{R_i^2}g\left(\frac{4}{3}\pi\rho\frac{x_i^3}{R_i}\right) + \frac{1}{4a_i^2 R_i^2} \left[g\left(\frac{m_i}{R_i}\right) - g\left(\frac{m_{i-1}}{R_i}\right) \right]^2. \quad (7)$$

The last radiation like term in eq.(7) can be neglected in the large N limit [1, 2]. Namely, defining

$$x_i = i\Delta x, \quad N\Delta x = \sqrt{\frac{3}{8\pi\rho}},$$

where N denotes the number of shells one finds

$$m_{2n} - m_{2n-1} = \frac{n^2}{N^3} \sqrt{\frac{108}{8\pi\rho}},$$

$$m_{2n+1} - m_{2n} = \frac{6n^2 + 6n + 1}{N^3} \sqrt{\frac{3}{8\pi\rho}}.$$

So that for large enough values of N the difference $g(m_i/R_i) - g(m_{i-1}/R_i)$ becomes quite small that makes it possible to omit this term in the large N limit. Neglecting the last radiation like term and introducing a new function λ [7]

$$2g\left(\frac{2m}{r}\right) \equiv r^2 H_0^2 \lambda\left(\frac{2m}{r^3 H_0^2}\right),$$

from eq.(6) one finds

$$\left(\frac{\dot{R}_i}{R_i}\right)^2 = -k\left(\frac{x_i}{R_i}\right)^2 + H_0^2 \lambda\left(\frac{8\pi\rho}{3H_0^2} \frac{x_i^3}{R_i^3}\right), \quad (8)$$

which after redefinition $x_i/R_i \rightarrow \text{scale factor}$ determines the cosmology with an arbitrary curvature term corresponding to the modified gravity [6, 7].

To summarize we presented exact solution of dust shell universe dynamics in the case of modified gravity and by means of this solution derived the equation governing the cosmological evolution. Using the present solution one can continue in the spirit of papers [1, 2] to study an impact of small-scale inhomogeneities on the modified cosmology. However, one has to keep in mind that this model is highly idealized. In general local inhomogeneities strongly affect the light propagation, giving rise to dispersions in the observed distance-redshift relation.

Acknowledgments

The author is indebted to M. Gogberashvili for useful conversations. The work was supported by the grant REL.RIG.9807.

-
- [1] N. Sugiura, K. Nakao and T. Harada, Phys. Rev. **D58** (1998) 103504-1; gr-qc/9808012.
[2] N. Sugiura, K. Nakao and T. Harada, Phys. Rev. **D60** (1999) 103508-1; gr-qc/9911090.
[3] W. Israel, Nuovo Cim. **B44** (1966) 1; Erratum-ibid. **B48**(1967) 463.
[4] M. Milgrom, Astrophys. J. **270**, (1983) 365; M. Milgrom, Astrophys. J. **270**, (1983) 371; M. Milgrom, Astrophys. J.

- 270**, (1983) 384.
[5] R. Sanders and S. McGaugh, Ann. Rev. Astron. Astrophys. **40** (2002) 263; astro-ph/0204521.
[6] A. Lue, R. Scoccimarro and G. Starkman, Phys. Rev. **D69** (2004) 044005; astro-ph/0307034.
[7] A. Lue and G. Starkman, Phys. Rev. Lett. **92** (2004) 131102; astro-ph/0310005.